

A simultaneous journal / wiki publication and dissemination of a new species description: *Neobidessodes darwiniensis* sp. n. from northern Australia (Coleoptera, Dytiscidae, Bidessini)

Lars Hendrich^{1,†}, Michael Balke^{1,2,‡}

1 Zoologische Staatssammlung, Münchhausenstrasse 21, D-81247 München, Germany **2** GeoBio Center, Ludwigs-Maximilians-Universität, München, Germany

† urn:lsid:zoobank.org:author:06907F16-4F27-44BA-953F-513457C85DBF

‡ urn:lsid:zoobank.org:author:945480F8-C4E7-41F4-A637-7F43CCF84D40

Corresponding author: Lars Hendrich (hendrich@zsm.mwn.de)

Academic editor: Martin Fikáček | Received 15 December 2010 | Accepted 2 February 2011 | Published 3 February 2011

urn:lsid:zoobank.org:pub:41390A60-77D6-457D-BC1C-6841717F6B21

Citation: Hendrich L, Balke M (2011) A simultaneous journal / wiki publication and dissemination of a new species description: *Neobidessodes darwiniensis* sp. n. from northern Australia (Coleoptera, Dytiscidae, Bidessini). ZooKeys 79: 11–20. doi: 10.3897/zookeys.79.803

Abstract

Here, we describe a new Australian species in journal format and simultaneously open the description in a wiki format on the www.species-id.net. The wiki format will always link to the fixed original journal description of the taxon, however it permits future edits and additions to species' taxonomy and biology. The diving beetle *Neobidessodes darwiniensis* **sp. n.** (Coleoptera: Dytiscidae, Bidessini) is described based on a single female, collected in a rest pool of the Harriet Creek in the Darwin Area, Northern Territory. Within *Neobidessodes* the new species is well characterized by its elongate oval body with rounded sides, short and stout segments of antennae, length of body and dorsal surface coloration. In addition to external morphology, we used mitochondrial *cox1* sequence data to support generic assignment and to delineate the new species from other Australian Bidessini including all other known *Neobidessodes*. Illustrations based on digital images are provided here and as online resources. A modified key is provided. Altogether ten species of the genus are now known worldwide, nine from Australia and one from New Guinea.

Keywords

Wiki, species ID, online species pages, *cox1* sequence data, DNA barcoding, molecular biodiversity assessment

Introduction

Many approaches and initiatives to “accelerate” the descriptive taxonomic process have recently been proposed or partially implemented. We suggest that the wikimedia engine provides one of the most powerful tools for routine taxonomic work, with wikipedia providing generic data and wikispecies a taxonomic backbone, i.e. the tree of life (see Page 2010). Here, we test an approach where we publish a new species in open-access journal format, and at the same time upload the data to a purpose-built wiki, the species ID site, flanked by a wikispecies entry which *de facto* serves as a “shop window”.

The epigeal species of the Australasian genus *Neobidessodes* Hendrich & Balke, 2009 were recently treated in a comprehensive systematic revision, including morphological and molecular data (Hendrich et al. 2009). Two new species, one from Australia and one from New Guinea, were described. Larvae of the genus were described in Michat et al. (2010). In northern Australia, *Neobidessodes* are among the most common and widespread diving beetles occurring in rest pools of intermittent streams during the dry season. Despite the fact that the first author studied more than 6000 specimens from his own samples and numerous museum collections (Hendrich et al. 2009), the new species described in this publication is known just from the female holotype. The single specimen was until recently overlooked in a vial, including numerous *Neobidessodes mjobergi* (Zimmermann 1922) and *Hydroglaphys godeffroyi* (Sharp 1882) collected in August 2006, on the way from Pine Creek to the Kakadu National Park.

Combining morphology and mitochondrial DNA sequence data we describe the new species and provide a modified key for all epigeal species of the genus. The DNA sequence data and a high resolution digital image of the beetle habitus, coloration and sculpture are made available online for faster dissemination of taxonomic knowledge. Links are provided below.

Material and methods

Material. This study is based on the examination of 26 specimens, the holotype of our new species and specimens of *Neobidessodes bilita* (Watts, 1978) and *N. mjobergi*, deposited in CLH, SAMA and ZSM.

Neobidessodes bilita (Watts, 1978): Australia, New South Wales. 12 exs., S NSW, 6.5 km SW Eden, Towamba Road 2 km N Nullica, 556 m, 16.XI.2006, 37.04.412S 149.51.200E, L. & E. Hendrich leg. (NSW 111), two specimens with “DNA M.Balke 1900”, “DNA M.Balke 1901” [green printed labels] (CLH, ZSM).

Neobidessodes mjobergi (Zimmermann, 1922): Australia, Northern Territory. 13 exs., Manton Dam Recreation Area, 46 km S Darwin, 35 m, 19.VIII.2006, 12.50.270S 131.08.050E, L. & E. Hendrich leg. (NT 1), one specimen with “DNA M.Balke 1656” [green printed label] (CLH, ZSM).

Descriptions. Beetles were studied with a Leica MZ 12.5 dissecting scope at 10–100x. Habitus photos of beetles were made by Alexander Riedel (Karlsruhe, Germany) and by the authors. Image stacks were aligned and assembled with the computer software Helicon Focus 4.77TM. Abbreviations used in the text are: TL (total length), TL-H (total length without head), and MW (maximum width). Label data of type material are cited in quotation marks.

DNA sequencing and data analysis. We extracted DNA from the alcohol preserved female holotype after removal of the abdomen, using the Qiagen Dneasy tissue kit. We ran a PCR with Bioline Mago Taq at 94° for 2 min, 40 cycles of 94° for 30 s, 47° for 30 s and 72° for 60 s, and a final extension of 72° for 10 min, using primers for the 3' end of cytochrome c oxidase 1 (*cox1*) Jerry (F: 5'- CAA CAT TTA TTT TGA TTT TTT GG -3') and Pat (R: 5'- TCC AAT GCA CTA ATC TGC CAT ATT A -3') (Simon et al. 1994).

This *cox1* fragment is our standard “DNA barcoding” fragment for Dytiscidae, a short fragment of DNA used for preliminary species identification and study of population-level processes (see e.g. Hendrich and Balke 2009 and Hendrich et al. 2009). We used the *cox1* fragment sequenced for the female holotype to obtain quantitative data for species recognition which we here suggest especially useful as we had no male specimens for study of male genital structures. The sequence was added to our database of Australian Dytiscidae (Hendrich et al. 2010), containing around 70% of the Australian fauna, including all *Neobidessodes* species (Hendrich et al. 2009). We ran neighbour joining analysis in PAUP* (Swofford 2002) using HKY85 as well as uncorrected p-distances. The Species Identifier module of Taxon DNA software was used to study sequence divergence in the dataset (Meier et al. 2006).

Codens

CLH Collection Lars Hendrich, Berlin, Germany; property of the Naturhistorisches Museum Wien, Austria

SAMA South Australian Museum, Adelaide, South Australia, Australia

ZSM Zoologische Staatssammlung München, Munich, Germany

Results and discussion

DNA Sequencing

We obtained 450 bp 3' *cox 1* sequence (GenBank accession # FR733592). Ran against our 1400+ Australian *cox1* sequence database, we find minimum uncorrected p-distances in SpeciesIdentifier of 10.15% (*Limbodessus jundeensis* Watts and Humphreys, 2003), followed by *Neobidessodes samkrisi* Hendrich & Balke, 2009, *N. thoracicus* Hendrich &

Balke, 2009 and *N. bilita* (Watts, 1978) (10.37–10.39%) and e.g. *Copelatus tenebrosus* Régimbart, 1880 (10.59%). The neighbour joining analysis in PAUP* placed *N. darwiniensis* sp. n. as the sister of all other *Neobidessodes*. This is not necessarily the correct phylogenetic position of *N. darwiniensis*, but indicates that the female studied here does not belong to any other known *Neobidessodes*, nor to any other species in our database.

Taxonomy

Neobidessodes is a genus with 10 species distributed in Australia (9 species) and New Guinea (1 species). All but two [the stygobitic *N. limestonensis* (Watts and Humphreys, 2003) and *N. gutteridgei* (Watts & Humphreys, 2003)] species have a more or less contrasting black/yellow surface. The basic pattern of these species includes various yellow or reddish spots. The median lobes are simple and very elongate, in ventral view strongly tapered or rounded at tip. The size of the species varies from 1.95 to 3.85 mm (see also Hendrich et al. 2009).

The new species was placed in *Neobidessodes* because of the following combination of characters: 1) body elongate oval; 2) basal pronotal striae sharply incised, not connected by a transverse groove; 3) elytra lacking basal striae and sutural striae; 4) epipleura lacking transverse carina; 5) head lacking cervical line and its foremargin not bordered; 6) prosternal process broad, distinctly excavated and margined; 7) inner margin of both metacoxal wings strongly ridged; 8) hind margin of abdominal ventrites 3–5 without row of minor irregular dentate processes.

Checklist of *Neobidessodes* species

NSW = New South Wales; **NT** = Northern Territory; **QLD** = Queensland; **VIC** = Victoria; **WA** = Western Australia; **N** = northern; **S** = southern.

Australia – epigeal

<i>N. bilita</i> (Watts, 1978)	S QLD, NSW, VIC
<i>N. darwiniensis</i> Hendrich & Balke, sp. n.	NT
<i>N. denticulatus</i> (Sharp, 1882)	N WA, NT, QLD, NSW
<i>N. flavosignatus</i> (Zimmermann, 1922)	N WA, NT, N QLD
<i>N. grossus</i> (Zimmermann, 1922)	N WA, NT, N QLD
<i>N. mjobergi</i> (Zimmermann, 1922)	N WA, NT, N QLD
<i>N. thoracicus</i> Hendrich & Balke, 2009	N WA, NT, N QLD

Australia – stygobitic

<i>N. gutteridgei</i> (Watts & Humphreys, 2003)	WA (Yilgarn)
<i>N. limestonensis</i> (Watts & Humphreys, 2003)	WA (Yilgarn)

New Guinea – epigeal

<i>N. samkrisi</i> Hendrich & Balke, 2009	West Papua, Merauke, Indonesia
---	--------------------------------

***Neobidessodes darwiniensis* sp. n.**

urn:lsid:zoobank.org:act:CAD876B9-A027-458D-88D1-47E4A239FFA7

http://www.species-id.net/wiki/Neobidessodes_darwiniensis

Figs 1, 5, 6, 7

Type locality. Rest pool, Harriet Creek at Kakadu Highway, 11 km NE Pine Creek, Northern Territory, Australia [13°45'04.63"S 131°53.55.31"E].

Type material. Holotype: Female, "Australia: NT, Kakadu Hwy, Harriet Creek at Hwy Cross., 156m, 24.VIII.2006, 13.74.4816S 131.89.7483E, L. & E. Hendrich leg. (NT 14)"; "DNA M. Balke 3821" [green printed label]; "HOLOTYPE *Neobidessodes darwiniensis* sp. n. Hendrich & Balke 2010" [red printed label] (SAMA).

Description. Measurements. TL = 1.95 mm, TL-H = 1.8 mm; MW = 1.0 mm.


Colour. Antennae, palpi, head and most parts of pronotum reddish-brown, posterior angles of head, near eyes and base of pronotum in middle broadly dark brown. Elytron dark brown with some small vague yellow spots subbasally and subapically (Fig. 1). Ventral side, including legs and epipleura, reddish-brown, prosternal process and metacoxal plates somewhat darkened.

Sculpture and structure. Elongate oval, sides well rounded. Maximum width at apical third of body. Segments of antennae short and stout. Head with relatively coarse punctures and strong microreticulation. Pronotum and elytron with rather dense, medium-sized punctures and weak to moderate microreticulation, finely pubescent. Pronotal striae deep and well marked, length almost 1/2 of that of pronotum, strongly incurved converging anteriorly (Fig. 1). Elytra lacking basal and sutural striae. Under-side with a few moderately large weak punctures at sides, midline of metaventrite with moderately dense smaller punctures. Metacoxal lines raised, well separated, weakly diverging anteriorly.

Male. Unknown.

Female. Pro- and mesotarsi simple. Inner edge of mesotibia nearly straight.

Affinities - DNA Sequence Data. The 3' *cox1* sequence available at <http://www.ncbi.nlm.nih.gov/nuccore/FR733592.1> indicates that the new species is rather distinctive, the closest uncorrected p-distances in our database were other *Neobidessodes* species (c. 10.37%) and *Limbodessus jundeensis* (10.15%). **Morphology.** The smallest species of the genus. On first view, the new species resembles in size and colour the common *Hydroglyphus godeffroyi* (Fig. 4) distributed all over northern Australia and New Caledonia, and can be easily overlooked in the field. When recognized as a *Neobidessodes* the new species is similar to *N. mjobergi* (Fig. 3) in coloration and to *N. bilita* (Watts, 1978) (Fig. 2) in size. From *N. mjobergi* it can be separated by its more broadly oval body, the much smaller size (*N. mjobergi* 2.55-2.65 mm) and unicolourous head, and from *N. bilita* by the darker dorsal surface, the short and stout segments of antennae, the rounded, broadly oval body, and the unflanged subapical part of the elytra (Figs 1, 2). Furthermore, *N. bilita* is a strictly south-eastern species with a disjunct distribution from southern Queensland to Victoria (Hendrich et al. 2009).




Figures 1–4. Habitus of **1** *Neobidessodes darwiniensis* sp. n. (holotype, female) **2** *N. bilita* (female) **3** *N. mjobergi* and **4** *Hydroglyphus godeffroyi* (scale bar = 2.0 mm) (Photos: M. Balke, A. Riedel).

Etymology. Named after the Darwin area in the Northern Territory; the specific epithet is an adjective in the nominative singular.

Distribution. Only known from the type locality at Harriet Creek, 11 km NE Pine Creek but probably more widespread in the Northern Territory (Fig. 5).

Habitat. The single specimen was collected in one of the rest pools of a rocky creek, with gloomy water and at least partly shaded by smaller gum trees. The bottom consisted of coarse sand with a thick layer of unrotten leaves and twigs, no submerged or emergent vegetation visible (Figs 6, 7).

Neobidessodes darwiniensis sp. n. was associated with the dytiscids *Clypeodytes larsoni* Hendrich & Wang, 2006, *Hydroglyphus daemeli* (Sharp, 1882), *H. godeffroyi*, *H. grammopterus* (Zimmermann, 1928), *Hyphydrus contiguus* Wehncke, 1877, *H. lyra-tus* Swartz, 1808, *Laccophilus cingulatus* Sharp, 1882, *L. sharpi* Régimbart, 1889, *L.*



Figures 5–7. 5 Distribution of *Neobidessodes darwiniensis* sp. n. in Northern Australia. 6–7 Habitat of *Neobidessodes darwiniensis* sp. n., *N. grossus*, *N. mjobergi* and *N. thoracicus*, Northern Territory Kakadu Hwy, Harriet Creek at Hwy Crossing (NT 14) (Photos: L. Hendrich).

walkeri J. Balfour-Browne, 1939, *Limbodessus compactus* (Clark, 1862), *Neobidessodes grossus* (Zimmermann, 1922), *N. mjobergi*, *N. thoracicus* Hendrich & Balke, 2009, *Sternopriscus alligatorensis* Hendrich & Watts, 2004, *S. aquilonaris* Hendrich & Watts, 2004, *Tiporus centralis* (Watts, 1978), *T. guiliani* (Watts, 1978) and *T. undecimmaculatus* (Clark, 1862).

Remarks. Despite the fact that thousands of *Neobidessodes* were collected on three field trips to the Northern Territory and the Kimberley region, surprisingly only one specimen of *N. darwiniensis* sp. n. appeared. Most of the expeditions took place during the dry period, between June and October, when most of the other species of the genus dominate the remaining rest pools and swamps. We assume the new species is more common in or just after the rainy season, from November to April, as was observed for *N. grossus* (Hendrich et al. 2009).

The key to epigeal species of *Neobidessodes* in Hendrich et al. (2009) should be modified as follows:

- 1 Length > 3.7 mm. Elytron with a subapical lateral flange, pronotal striae very weak, N WA, NT, N QLD ***grossus***
- Length < 3.7 mm **2**
- 2 Elytron with a subapical lateral tooth, pronotal striae well marked, WA, NT, QLD, N NSW ***denticulatus***
- Elytron lacking lateral tooth, pronotal striae present or absent **3**
- 3 Pronotal striae absent **4**
- Pronotal striae present **5**
- 4 Length 2.55–2.65 mm, outline of junction of pronotum and elytra smooth, sides of pronotum evenly curved, maximum width at posterior angles, dorsal colour pattern usually diffuse, N WA, NT, N QLD ***mjobergi***
- Length 2.75–2.9 mm, outline of junction of pronotum and elytra slightly sinuate, maximum width of pronotum somewhat before base. Dorsal colour pattern strongly varying, when present, usually well marked. In some specimens pronotum yellow, in others pronotum and elytra all black, N WA, NT, N QLD ***thoracicus***
- 5 Dorsal colour pattern diffuse. Pronotal striae well marked and long (1/4 to 1/3 of length of pronotum) **6**
- Contrasting yellowish markings on black elytra. Pronotal striae only slightly marked and short (maximum 1/4 of length of pronotum) **7**
- 6 Body elongate oval. Pronotum as broad as elytra, outline of junction of pronotum and elytra slightly sinuate, maximum width of pronotum somewhat before base (Fig. 2). Males with mesotibia curved, length 2.2–2.25 mm, VIC, NSW, S QLD ***bilita***
- Body broader oval. Pronotum narrower than elytra, outline of junction of pronotum and elytra smooth, sides of pronotum evenly curved, maximum

- width at posterior angles (Fig. 1). Male unknown, length 1.95 mm, smallest species of the genus, NT ***darwiniensis* sp. n.**
- 7 Males with mesotarsus straight. Pronotal striae well marked but short, small species, length 2.0 mm, West Papua, Indonesia..... ***samkrisi***
- Males with mesotarsus straight. Pronotal striae extremely weak and faint, larger species, length 2.35–2.65 mm, N WA, NT, N QLD..... ***flavosignatus***

Acknowledgements

We are indebted to Hans Fery (Berlin, Germany), Chris Watts (Adelaide, Australia) and one additional, anonymous reviewer for improving an earlier version of the manuscript. We thank Gregor Hagedorn (Berlin, Germany) for stimulating discussions on species ID and the wiki format and the ZooKeys team for helping to implement our journal publication / wiki upload approach. The Parks and Wildlife Commission of the Northern Territory, the Australian National Parks and Wildlife Service in Darwin and the Kakadu National Park are thanked for giving permission to conduct scientific research in the Northern Territory (Permit Numbers: 23929 and RK- 400/ RK- 660). This work was supported by grants to Michael Balke: Deutsche Forschungsgemeinschaft (BA 2152/4-1, 6-1 and 7-1), and to Lars Hendrich: Deutsche Forschungsgemeinschaft (HE 5729/1-1).

References

- Hendrich L, Balke M (2009) *Kakadudessus tomweiri*, a new genus and species of diving beetle from tropical northern Australia, based on molecular, phylogenetic and morphological data (Coleoptera, Dytiscidae, Bidessini). *Zootaxa* 2134: 49–59.
- Hendrich L, Hawlitschek O, Balke M (2009) The epigeal Australasian species of *Neobidessodes* gen.n. diving beetles—a revision integrating morphology, cybertaxonomy, DNA taxonomy and phylogeny (Coleoptera: Dytiscidae, Bidessini). *Zootaxa* 2288: 1–41.
- Hendrich L, Pons J, Ribera I, Balke M (2010): Mitochondrial *cox1* sequence data reliably uncover patterns of insect diversity but suffer from high lineage-idiosyncratic error rates. *Plos ONE* 5(12): e14448. doi:10.1371/journal.pone.0014448
- Meier R, Kwong S, Vaidya G, Ng PKL (2006) DNA Barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Systematic Biology* 55: 715–728.
- Michat MC, Alarie Y, Watts CHS (2010) Descriptions of the first-instar larva of the hypogaic species *Neobidessodes limestoneensis* (Watts & Humphreys) and of the third-instar larva of *Hydroglyphus balkei* Hendrich (Coleoptera: Dytiscidae: Bidessini) with phylogenetic considerations. *Zootaxa* 2658: 38–50.
- Page RDM (2010) Wikipedia as an encyclopaedia of life. *Organisms Diversity and Evolution* 10: 343–349.

- Simon C, Frati F, Beckenbach AT, Crespi B, Liu H, Flook P (1994) Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America* 87: 651–701.
- Swofford DL (2002) PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Sinauer Associates, Sunderland MA.